

Combinatorics for Metrical Feet

Abstract. This squib provides a proof that the number of parsings into metrical feet (Hayes 1995) of a string of length n without regard for headedness is the Fibonacci number $\text{Fib}(2n)$.

Key words: Fibonacci, combinatorics, metrical feet

1. Introduction

Halle and Vergnaud (1987) propose a convention on the parsing of elements into metrical feet, the *Exhaustivity Condition*, that requires all elements to belong to some foot, except for certain principled cases of *extrametricality*. However, the general consensus since Halle and Vergnaud (1987) is that metrical elements can remain unparsed, failing to belong to any foot, generalizing the notion of extrametricality. Hayes (1995), Halle and Idsardi (1995), and Kager (1999) among many others explicitly reject the Exhaustivity Condition. Hayes's comments are given in (1), Halle and Idsardi's are given in (2).

- (1) “The upshot seems to be that in our present state of knowledge, it would be aprioristic to adhere firmly to a rigid principle of exhaustive prosodic parsing ...”
Hayes 1995: 110.
- (2) “We also deviate from previous metrical theories by not requiring exhaustive parsing of the sequence of elements, that is we do not require that every element belong to some constituent ...” Halle and Idsardi 1995: 440.

In this squib I will prove that the number of possible metrical parsings into feet under these assumptions for a string of n elements is $\text{Fib}(2n)$ where $\text{Fib}(n)$ is the n^{th} Fibonacci number.

2. Initial observations

Disregarding prominence relations within the feet (i.e. headedness), the possible footings for strings up to a length of three elements are shown in (3). Feet are indicated by matching parentheses; elements not contained within parentheses are unfooted (*unparsed* in Optimality Theory terminology).

- (3) a. 1 element, 2 possible parsings: (x), x
 b. 2 elements, 5 possible parsings: (xx), (x)(x), (x)x, x(x), xx
 c. 3 elements, 13 possible parsings: (xxx), (xx)(x), (xx)x, (x)(xx), x(xx), (x)(x)(x),
 (x)(x)x, (x)x(x), x(x)(x), (x)xx, x(x)x, xx(x), xxx

The number of possible footings is equal to every other member of the Fibonacci sequence, defined and illustrated in (4); see for example Cameron (1994).

- (4) Fibonacci sequence

$$Fib(0) = Fib(1) = 1; \text{ for } n > 1 \text{ } Fib(n) = Fib(n-1) + Fib(n-2)$$

1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, 233, ...

There is only one possible footing of a string of zero elements, so that it is also the case that $footings(0) = Fib(0)$.

3. Proof

Let $footings(n)$ be the number of parsings of a string of n elements into metrical feet, not subject to the Exhaustivity Condition. We can derive a recurrence relation for the number of metrical feet in a string of length $n+1$ by dividing the string after the places where an initial foot could occur, as shown in (5).

- (3) a. no initial foot: x | ..., n elements left, therefore $footings(n)$ footings

b. 1 element foot: (x) | ..., n elements left, therefore $footings(n)$ footings

c. 2 element foot: (xx) | ..., n-1 elements left, therefore $footings(n-1)$ footings

d. 3 element foot: (xxx) | ..., n-2 elements left, therefore $footings(n-2)$ footings

...

e. n element foot (x...x)|, 0 elements left, therefore $footings(0) = 1$ footing

generally then, $footings(n+1) = footings(n) + \sum_{i=0}^n footings(i)$

We then prove the general relation by induction on n. That is, assuming that $footings(n) = Fib(2n)$, we will prove that $footings(n+1) = Fib(2n + 2)$. We begin with the recurrence relation derived in (3), and expand out the first term of the summation, as shown in (4).

$$(4) \quad footings(n+1) = footings(n) + footings(n) + \sum_{i=0}^{n-1} footings(i)$$

Substituting for $footings(n)$ using the induction assumption gives (5).

$$(5) \quad footings(n+1) = Fib(2n) + Fib(2n) + \sum_{i=0}^{n-1} footings(i)$$

Substituting for $Fib(2n)$ using the Fibonacci recurrence relation gives (6).

$$(6) \quad footings(n+1) = Fib(2n) + Fib(2n-1) + Fib(2n-2) + \sum_{i=0}^{n-1} footings(i)$$

Substituting for $Fib(2n-2)$ again using the induction assumption gives (7).

$$(7) \quad footings(n+1) = Fib(2n) + Fib(2n-1) + footings(n-1) + \sum_{i=0}^{n-1} footings(i)$$

Substituting for the last two terms using the footings recurrence relation gives (8).

$$(8) \quad footings(n+1) = Fib(2n) + Fib(2n-1) + footings(n)$$

Substituting for $footings(n)$ again using the induction assumption gives (9).

$$(9) \quad \text{footings}(n+1) = \text{Fib}(2n) + \text{Fib}(2n-1) + \text{Fib}(2n)$$

Substituting for the first two terms using the Fibonacci recurrence relation gives (10).

$$(10) \quad \text{footings}(n+1) = \text{Fib}(2n+1) + \text{Fib}(2n)$$

Again substituting using the Fibonacci recurrence relation gives (11), as required.

$$(11) \quad \text{footings}(n+1) = \text{Fib}(2n+2) \text{ Q.E.D.}$$

Having proved that if $\text{footings}(n) = \text{Fib}(2n)$ then $\text{footings}(n+1) = \text{Fib}(2n+2)$ for $n > 1$, and having $\text{footings}(0) = \text{Fib}(0)$ and $\text{footings}(1) = \text{Fib}(2)$, we have proved the relation for all non-negative n .

4. Corollary

Given the above proof, substituting into the footing recurrence relation gives (12).

$$(12) \quad \text{footings}(n+1) = \text{footings}(n) + \sum_{i=0}^n \text{footings}(i)$$

$$\text{Fib}(2n+2) = \text{Fib}(2n) + \sum_{i=0}^n \text{Fib}(2i)$$

And, since from the Fibonacci recurrence relation $\text{Fib}(2n+2) = \text{Fib}(2n+1) + \text{Fib}(2n)$, therefore we derive (13).

$$(13) \quad \text{Fib}(2n-1) = \sum_{i=0}^n \text{Fib}(2i)$$

That is, for example, $\text{Fib}(7) = \text{Fib}(6) + \text{Fib}(4) + \text{Fib}(2) + \text{Fib}(0) = 13 + 5 + 2 + 1 = 21$.

5. Conclusion

The number of non-exhaustive parsings of n elements into metrical feet (i.e. the number of non-exhaustive partitions of n elements) has been proven to be equal to $\text{Fib}(2n)$, the $2n^{\text{th}}$ Fibonacci number.

References

- Cameron, P. J. (1994) *Combinatorics: Topics, Techniques, Algorithms*. Cambridge University Press.
- Halle, M. and W. J. Idsardi (1995) General Properties of Stress and Metrical Structure. In Goldsmith, J. (ed.), *The Handbook of Phonological Theory*, Blackwell, chapter 11, pp. 403-443.
- Halle, M. and J.-R. Vergnaud (1987) *An Essay on Stress*. MIT Press.
- Hayes, B. (1995) *Metrical Stress Theory: Principles and Case Studies*. University of Chicago Press.
- Kager, R. (1999) *Optimality Theory*. Cambridge University Press.